□□□□□□□□□□

□□□□□□□□□□□□□□(□□□□□□□)

Image

□□□□□□□□□□□□STS□□

□□

□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ □□□□□ □□ □□ □□ □□

□□□□□□□□□□

□□※□□
□□□□□□□□□□□□□□□[1□]□

※□2002□□WHO□□□□□□□□□□□□□□□□MFH□□□□□□□□□□□UPS□/MFH□□□□□□□□□□□□□□□□□□2013□
□WHO□□□□□UPS/MFH□□□□/□□□□□□□undifferentiated/unclassified sarcoma□□□
□undifferentiated pleomorphic sarcomas□□□□□□

Image


悪性軟部腫瘍の組織型別頻度
1.2%
1.2%
1.5%
1.6%
1.6%
2.1%
2.4%
1.0%
0.7%
1.8%
39.4%
20.5%
8.0%
6.8%
5.6%
4.9%
脂肪肉腫
悪性線維性組織球腫
粘液線維肉腫
平滑筋肉腫
滑膜肉腫
悪性末梢神経鞘腫瘍
横紋筋肉腫
線維肉腫
骨外性軟骨肉腫
血管肉腫
骨外性ユーイング肉腫・未熟神経外胚葉性腫瘍
低悪性度線維粘液性肉腫
類上皮肉腫
胞巣状軟部肉腫
明細胞肉腫
その他の悪性軟部腫瘍
1,540 例
日本整形外科学会　骨・軟部腫瘍委員会／国立がん研究センター（編）.：全国軟部腫瘍登録一覧表（2012）

[illegible]

[illegible][2]

[illegible][2]

Image



対象・方法：SEER(Surveillance, Epide mi ology, and End Results) の登録データから、1998 ～ 2001 年の米国における悪性軟部腫瘍 28,758 例（20 歳以上）の組織型別生存率を評価した。

Ries LAG, Young JL, Keel GE, Eisner MP, Lin YD, Horner M-J (editors).
SEER Survival Monograph: Cancer Survival Among Adults: U.S. SEER Program, 1988-2001,
Patient and Tumor Characteristics. National Cancer Institute,
SEER Program, NIH Pub. No. 07-6215, Bethesda, MD, 89-92 (2007)

Image



対象・方法：MSKCC(Memorial Sloan Kettering Cancer Center) にて 1982 年 7 月～2001 年 6 月末までに治療を受けた悪性軟部腫瘍患者 1,097 例の生存率を評価した。

Edge S, Byrd DR, Compton CC, Fritz AG, Greene FL, Trotti A (editors). AJCC Cancer Staging Manual, 7th ed, Springer, 291-298 (2009)

1□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□.□□□□□□□□□□□□ (2012)
2□□□.□□□□□□□□□□□□□ 2012.□□□ (2012)
3□Ries LAG, Young JL, Keel GE, Eisner MP, Lin YD, Horner M-J (editors). SEER Survival Monograph: Cancer Survival Among Adults: U.S. SEER Program, 1988-2001, Patient and Tumor Characteristics. National Cancer Institute, SEER Program, NIH Pub. No. 07-6215, Bethesda, MD, 89-92 (2007)
4□Edge S, Byrd DR, Compton CC, Fritz AG, Greene FL, Trotti A (editors). AJCC Cancer Staging Manual, 7th ed, Springer, 291-298 (2009)

□□□□□□□□□□□□(STS)□□□□□□

□□□□□□□□□□□□□□□□□

□□□□□□

Source URL: *https://www.loadtest.pro.novartis.com/products/votrient/sarcoma/sts_feature*