NOVARTIS

□□□□□□□□□□□□

□□□□□□□□□□□□

□□□□□□

1.□□□□□

□□□□□□□□□□□□□□□□□□□□□□1□75□600mg□2□□4□□□□□□□□□□□□□1□□□□□□□□□□□□□□□□□□□□□□□□□□□
□IgE□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□

□□□□□□□1□□□□□

4□□□□□

Image

投与前の血清中総 IgE濃度(IU/mL)	体重(kg)									
	≧20～25	＞25～30	＞30～40	＞40～50	＞50～60	＞60～70	＞70～80	＞80～90	>90~125	>125~150
≧30～100	75mg	75mg	75mg	150mg	150mg	150mg	150mg	150mg	300mg	300mg
＞100～200	150mg	150mg	150mg	300mg	300mg	300mg	300mg	300mg	450mg	600mg
＞200～300	150mg	150mg	225mg	300mg	300mg	450mg	450mg	450mg	600mg	
＞300～400	225mg	225mg	300mg	450mg	450mg	450mg	600mg	600mg		
＞400～500	225mg	300mg	450mg	450mg	600mg	600mg				
＞500～600	300mg	300mg	450mg	600mg	600mg					
＞600～700	300mg		450mg	600mg						
＞700～800										
＞800～900										
>900~1,000										
>1,000~1,100										
>1,100~1,200										
>1,200~1,300										
>1,300~1,500										

4週間毎投与の表に該当しない場合には
2週間毎投与の表に従い投与すること

2□□□□□

Image

投与前の血清中総IgE濃度(IU/mL)	体重(kg)									
	≥20～25	>25～30	>30～40	>40～50	>50～60	>60～70	>70～80	>80～90	>90～125	>125～150
≥30～100										
>100～200										
>200～300										375mg
>300～400									450mg	525mg
>400～500							375mg	375mg	525mg	600mg
>500～600						375mg	450mg	450mg	600mg	
>600～700		225mg			375mg	450mg	450mg	525mg		
>700～800	225mg	225mg	300mg	375mg	450mg	450mg	525mg	600mg		
>800～900	225mg	225mg	300mg	375mg	450mg	525mg	600mg			
>900～1,000	225mg	300mg	375mg	450mg	525mg	600mg				
>1,000～1,100	225mg	300mg	375mg	450mg	600mg					
>1,100～1,200	300mg	300mg	450mg	525mg	600mg					
>1,200～1,300	300mg	375mg	450mg	525mg						
>1,300～1,500	300mg	375mg	525mg	600mg						

2週間毎投与の表に該当しない場合には4週間毎投与の表に従い投与すること

投与不可

□□□□□□□□□□□□□□□□□□□□□□□□0.008mg/kg/□IU/mL□□□□2□□□□□□□□□□□□□□0.016mg/kg/□IU/mL□□□□4□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□

□□□□PD□

□□IgE□□□□□□□□□□□□□□□□□□

□□□□□□□IgE□□□□□□□

Image



2.□□□IV□□□□□□□□2210□□□□□□□□□□□□[1]□

1□ Zielen, S. et al.: Int. Arch. Allergy Immunol. 160□1□, 102, 2013□□XOLM01866□

□□□□□□□□□□

□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□5 □□□□□□□□□□□□□□□□□□□□□□□□□□□□□

□□□□□□□□□□□□□□□□□□□□□□IgE□□□1,500□2,000IU/mL□□□□□□□□□□□□□□□□□
□□
□□□□□

□1□□□□□

□□

□□□600mg□□□2□□□□□□□□□□□□□□□□ABP□□□□□□□□□□□□□EAR□□□□□□□□□□IgE□
□2,000IU/mL□□□□□□□□□□□□□□□□□□□□□

□□□

□□□□□□□□□□□□□□□□□□□□□□□

□□□

□□□□□□□□□ABP□□□□EAR□□□□□10□30□□FEV1.0□□□□□□□□□□□□□□PK□□□□□PD□□□□
□□□
□□□□□□□□ABP□□□□□□□□□□LAR□□□□□□□3□8□□□FEV1.0□□□□□□PK/PD□□□IgE□□□IgE
□□FENO□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□

□□

□□□□□□18□65□□□40□150kg□50□[□1]
□Group1□IgE□□30□300IU/mL□Group2□□700□2,000IU/mL□

□1 □□□□□□□□□□□□□□□□□□□□□

□□□

□□□□1□□□□□□□□□□□□□□□□□□□□□□□□□IgE□□□□□□□□□[□2]□□□□□□□□□□□□□□□□□
□□□□□□□□Group1□□□□□□□□□□□□□1□75□375mg□4□□2□□□□□□□□□Group2□□□□□1□□
□600mg□2□□□□□□□□□
□□□□□□□□IgE□□□□□□□□□□□□□□□2,000IU/mL□□□□□□□□□□□□□□1□600mg□2□□□□□□

□2 Group1□□□□□□2008□6□□EU□□□□□□□□□□□□□□□□□□□□
Group2□□□□□□□□□□□□□□□□IgE□□□11ng/mL□□□□□□□□□□□□□□□□□□□□□□□

□□□□

Image



□2□ABP□□□□□EAR□□□□□□□□□□□ □□□□□□ □

□□8□□□EAR□□□□□□□□23.1%□□□□□□□□□□□□□Group1□9.3%□Group2□5.6%□□□□□□□□□□□□□□□□□□□p=0.018□□p<0.001□ANCOVA□□□□16□□□EAR□□□□□□□□20.0%□□□□□□□□□□□□□Group1□11.8%□Group2□5.1%□□□□□p=0.087□□p<0.001□ANCOVA□□

4. □□□□□□□□□□□
□ □□

5. □□□□□□□□□□□□□□□□□□□
□□□□□□□
□□IgE□□
□□□□□
• □□□□□□□□□□□□□□□□□□□□FEV1.0□□□□□□□□□□80%□□□
• □□□□□□□□□□□□□
• □1□□□□□□□□□□□□□□

□3□□□□□□PK□

□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□

□□□□□□□□□□□□□□□□□□□□□

Image



[illegible]4[illegible]

[illegible]Group1[illegible]50.0%[illegible]9/18[illegible]Group2[illegible]100.0%[illegible]16/16[illegible]75.0%[illegible]12/16[illegible]Group1[illegible]3[illegible]16.7%[illegible]Group2[illegible]5[illegible]31.3%[illegible]4[illegible]25.0%[illegible]Group1[illegible]1[illegible]5.6%[illegible]Group2[illegible]1[illegible]6.3%[illegible]2.9%[illegible]1/34[illegible]6.3[illegible]1/16[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

PFS[illegible]

[illegible]

[illegible]

IC[illegible]

IC[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

IgE[illegible]

Image

Source URL: *https://www.loadtest.pro.novartis.com/products/xolair/asthma/capacity*